



FACT SHEET



Product:	Detoxin™
Suitable for:	Gently eliminating toxins from the body
Ingredients:	Cyanara Artichoke 100mg Dandelion root 100mg Nettle leaf 50mg Aloe vera extract 50mg Cranberry powder 50mg Green tea extract 25mg Barley grass extract 25mg Sea kelp 25mg Alfalfa leaf 25mg Parsley leaf 25mg Dicalcium phosphate Microcrystalline cellulose Magnesium stearate
Recommended dose:	For adults only – take 1 tablet a day. Do not exceed the stated dose.

- Contraindications: Detoxin™ is not recommended for people who are allergic to artichokes or other members of the daisy family, or those with gallstones. Detoxin™ should not be taken during pregnancy or whilst breastfeeding.
- Documented side effects: Nettle leaf may cause mild stomach upset in some people.
- Drug interactions: Drug interactions are possible between dandelion root and some diuretics and antibiotics and between cranberry and warfarin. Green tea may interact with some painkillers, some heart drugs, drugs used to treat the common cold and hayfever, some anti-diarrhoeal medications and warfarin.

Always tell your health care providers about any complementary and alternative practices you use. Providing a full picture of what you do to manage your health will help ensure coordinated and safe care.

Further reading:

Kirchhoff R, Beckers C, Kirchhoff G, Trinczek-Ga"rtner H, Petrowicz O, Reimann H. Increase in choleresis by means of artichoke extract. Results of a randomized placebo-controlled double blind study. *Phytomedicine* 1994; 1: 107–15.

Gebhardt R, Fausel M. Antioxidant and hepatoprotective effects of artichoke extracts and constituents in cultured rat hepatocytes. *Toxicol Vitro* 1997 ;11: 669–72.

Gebhardt R. Inhibition of hepatic cholesterol biosynthesis by artichoke leaf extracts is mainly due to luteolin. *Cell Biol Toxicol* 1997; 13: 58.

Brown JE, Rice-Evans CA. Luteolin-rich artichoke extract protects low density lipoprotein from oxidation in vitro. *Free Rad Res* 1998; 29: 247–55.

Pe´rez-Garcia F, Adzet T, Canigueral S. Activity of artichoke leaf extract on reactive oxygen species in human leukocytes. *Free Rad Res* 2000; 33: 661–5.

Kuusi T, Pyylaso H, Autio K. The bitterness properties of dandelion. II. Chemical investigations. *Lebensm-Wiss Technol* 1985;18:347–9.

Racz-Kotilla E, Racz G, Solomon A. The action of *Taraxacum officinale* extracts on body weight and diuresis of laboratory animals. *Planta Med* 1974;26:212–7.

Schütz K, Carle R, Schieber A. *Taraxacum*--a review on its phytochemical and pharmacological profile. *Ethnopharmacol.* 2006 Oct 11;107(3):313-23. Epub 2006 Jul 22.

Grieve M. *A Modern Herbal*, Vol 2. New York, NY: Dover Publications; 1971:574-579.

Odes HS, Madar Z. A double-blind trial of a celandin, aloe vera and psyllium laxative preparation in adult patients with constipation. *Digestion.* 1991;49(2):65-71.

Imai K, Nakachi K. Cross sectional study of effects of drinking green tea on cardiovascular and liver diseases. *BMJ* 1995;310:693-6.

Goto K, Kanaya S, Nishikawa T, et al. The influence of tea catechins on fecal flora of elderly residents in long-term care facilities. *Ann Long-Term Care* 1998;6:43-8.

Shirai N, Suzuki H. Effects of simultaneous intakes of fish oil and green tea extracts on plasma, glucose, insulin, C-peptide, and adiponectin and on liver lipid concentrations in mice fed low- and high-fat diets. *Ann Nutr Metab.* 2008;52(3):241-9. Epub 2008 Jun 19.

Jin X, Zheng RH, Li YM. Green tea consumption and liver disease: a systematic review. *Liver Int.* 2008 May 14. [Epub ahead of print]

Molgaard J, von Schenck H, Olsson AG. Alfalfa seeds lower low density lipoprotein cholesterol and apolipoprotein B concentrations in patients with type II hyperlipoproteinemia. *Atherosclerosis* 1987;65:173-9.

Behall KM, Scholfield DJ, Hallfrisch J. Diets containing barley significantly reduce lipids in mildly hypercholesterolemic men and women. *Am J Clin Nutr.* 2004 Nov;80(5):1185-93.